

# Mannitol

ALLEN R. NISSENSON, MD; RAYMOND E. WESTON, MD, and  
CHARLES R. KLEEMAN, MD, Los Angeles

*Mannitol may be useful clinically both as a diuretic and as an obligate extracellular solute. As a diuretic it can be used to treat patients with intractable edema states, to increase urine flow and flush out debris from the renal tubules in patients with acute tubular necrosis, and to increase toxin excretion in patients with barbiturate, salicylate or bromide intoxication. As an obligate extracellular solute it may be useful to ameliorate symptoms of the dialysis disequilibrium syndrome, to decrease cerebral edema following trauma or cerebrovascular accident, and to prevent cell swelling related to renal ischemia following cross-clamping of the aorta. Largely unexplored uses for mannitol include its use as an osmotic agent in place of dextrose in peritoneal dialysis solutions, its use to maintain urine output in patients newly begun on hemodialysis, and its use to limit infarct size following acute myocardial infarction.*

MANNITOL IS A 6-carbon alcohol with a molecular weight of 182, prepared commercially by the reduction of dextrose. Although it has an important place in current therapy, few physicians fully understand its chemistry, mode of action or clinical applicability. Since 1940, when Smith and associates<sup>1</sup> showed that mannitol clearance closely reflected glomerular filtration rate in man, there has been clinical interest in mannitol. However, recent emphasis on the role of cell swelling in the genesis of ischemic tissue damage has rekindled awareness of the medical uses of mannitol as more than only an osmotic diuretic.

From the Department of Medicine, UCLA School of Medicine (University of California, Los Angeles).

Submitted, revised, June 11, 1979.

Supported in part by a grant from McGaw Laboratories, Inc., Irvine, California.

Reprint requests to: Allen R. Nissenson, MD, Assistant Professor of Medicine, UCLA Hospital and Clinics, 10833 LeConte Ave., Los Angeles, CA 90024.

## Pharmacokinetics

Because orally administered mannitol is not absorbed, it must be administered parenterally. It distributes almost entirely in the extracellular fluid, very little penetrating cells. As a result, it is virtually inert, only 7 percent to 10 percent being metabolized, probably in the liver, while the rest is freely filtered by the glomeruli and excreted intact in the urine. About 7 percent is reabsorbed by the renal tubules. With normal kidney function, after a single intravenous dose, the half-life of mannitol in the circulating plasma is 15 minutes.<sup>2</sup> Of an injected dose, 90 percent is recovered in the urine after 24 hours.<sup>3</sup> However, with severe renal insufficiency the rate of mannitol excretion is greatly reduced and retained mannitol may increase extracellular tonicity leading to a shift of water out of cells, expanding the extracellular

fluid as well as inducing an apparent hyponatremia with increased serum osmolality such as occurs in hyperglycemia. Therefore, mannitol should be used cautiously under these conditions.

## Physiological Effects

### Renal

Osmotic diuretics are obligate solutes of low molecular weight, which are freely filtered at the glomerulus and poorly reabsorbed by the renal tubules. The resulting high urinary concentration profoundly affects renal water and sodium reabsorption. An almost ideal prototype, mannitol was used extensively in early physiological studies of osmotic diuresis. Wesson and Anslow<sup>4</sup> concluded that the proximal tubule was the primary site of action of mannitol's osmotic diuretic effect. More recently, however, Seely and Dirks<sup>5</sup> have precisely defined this phenomenon, their work being confirmed later by others.<sup>6</sup> Using micropuncture techniques, they elegantly demonstrated that half the effect of mannitol occurs proximally and half occurs in the ascending limb of the loop of Henle and proposed the following mechanism: The osmotic effect of mannitol depresses proximal reabsorption of water more than sodium, thereby decreasing proximal tubular fluid sodium concentration. Consequently, the gradient for passive sodium reabsorption in the thin ascending limb of the Henle loop is decreased. The natriuresis of mannitol diuresis, therefore, results largely from decreased sodium reabsorption in the loop of Henle. In addition, changes in medullary tonicity secondary to a mannitol augmentation of medullary blood flow may also effect sodium excretion.<sup>7</sup> As sodium delivery to distal tubular exchange sites increases, potassium excretion may rise during mannitol diuresis.

The effect of mannitol on renal hemodynamics may be both direct and indirect. Ternes and co-workers<sup>8</sup> showed that there was a dose dependent increase in renal vascular resistance in isolated perfused dog renal artery strips exposed to varying perfusate concentrations of mannitol. In contrast, others<sup>9-11</sup> found there to be an increase in total renal blood flow *in vivo* in animals, independent of the plasma volume increase. No comparable data for humans are available.

Lilien<sup>12</sup> reported a biphasic dose related response in rats, which might provide a unified view of the mannitol effect on renal hemody-

namics. At lower doses, there is definite vasodilatation probably mediated through efferent arteriolar dilatation, which could also contribute to the slight decrease in glomerular filtration rate during mannitol infusions. Conversely, at higher doses, the vasoconstrictor effect reported by Ternes and associates<sup>8</sup> supervenes.

Glomerular filtration rate falls slightly in animals and humans given mannitol. When mannitol is present in the proximal nephron and prevents, through its osmotic effect, sodium and water reabsorption, filtrate volume and pressure in this nephron segment rise. It is this increased intratubular pressure after mannitol administration that decreases glomerular filtration rate by Starling's law. Efferent arteriolar dilatation after mannitol is administered may occur (as has been mentioned) and may also play a role in the decreased glomerular filtration rate.

When proximal tubular sodium reabsorption is inhibited for any reason, calcium excretion in the urine increases. The precise mechanism of this coupling of sodium and calcium excretion is unknown. Even though, as explained above, decreased proximal tubular sodium reabsorption does not account for the bulk of the natriuresis after mannitol administration, the amount of sodium escaping proximal tubular reabsorption is increased. As a result, calcium excretion increases proportionately.<sup>13,14</sup> Independent of the natriuresis, urinary phosphate excretion also increases during mannitol diuresis.<sup>15</sup>

The only renal histologic effect of mannitol infusions in animals<sup>16</sup> or humans<sup>17</sup> is the same reversible proximal tubular cell vacuolization reported after sucrose and other osmotic diuretics. However, it is not associated with any functional tubular defects.

### Cardiovascular

Intravenously injected mannitol is confined to the extracellular fluid space. The increased tonicity obligates water to move from the intracellular to the extracellular fluid to maintain osmotic equilibrium. With each 182 mg per dl increase in mannitol concentration, a 10 milliosmole increase in extracellular fluid osmolality occurs with the same proportionate dilution of all serum electrolytes as occurs in hyperglycemia. Various amounts of plasma volume expansion have been reported with mannitol infusion ranging from 1 ml per kg of body weight per 5 grams of mannitol infused<sup>18</sup> to 3 ml per kg of body weight per 5 grams of

mannitol infused,<sup>19</sup> depending on body osmolarity and the relationship among plasma volume, extracellular fluid and total body water in the individual animal or person. With each 100 mg per dl increase in mannitol concentration in the extracellular fluid, serum sodium falls 1.6 to 2.6 mEq per liter.

After the expansion of the extracellular fluid and blood volume by injection of mannitol mean arterial pressure,<sup>20,21</sup> cardiac output,<sup>20,21</sup> heart rate,<sup>21,22</sup> coronary blood flow<sup>20</sup> and left ventricular end diastolic pressure<sup>22</sup> all increase. In isolated perfused artery segments, physiologic doses of mannitol cause vasodilatation and block the vasoconstrictor effects of norepinephrine.<sup>23</sup>

### *Cerebrovascular Disease*

Mannitol has several physiologic effects on the cerebrovascular system. Increased cerebral blood flow,<sup>24</sup> increased cerebral oxygen consumption<sup>24</sup> and decreased cerebrospinal fluid pressure<sup>24,25</sup> have all been documented. The decreased cerebrospinal fluid pressure, in spite of increased cerebral blood flow, is of interest. Since mannitol cannot cross the blood brain barrier, it causes water to move out of the brain into the extracellular fluid. This decrease in brain water is the cause of the fall in cerebrospinal fluid pressure. The movement of water out of the brain in response to the osmotic effect of mannitol decompresses the brain when cerebral edema is present. It is more effective than urea in this regard since urea slowly crosses the blood brain barrier and therefore maintains a lower osmotic gradient for water movement out of the brain.

### *Ocular System*

Mannitol profoundly lowers intraocular pressure when given intravenously<sup>26,27</sup> just as it lowers cerebrospinal fluid pressure. In addition, when applied topically to the cornea mannitol may be absorbed and also have an ocular hypotensive effect.<sup>28</sup>

### **Concept and Importance of Cell Swelling**

In a variety of organs, including kidney, heart and brain, following relatively brief periods of vascular occlusion, reperfusion fails to restore normal circulation.<sup>29,30</sup> This phenomenon of impaired reflow is caused by vascular endothelial cell swelling induced by the ischemic anoxia. Normally, the diffusion of sodium into and of potassium out of cells down existing concentra-

tion gradients is opposed by a *pump* requiring energy from oxidative metabolism, which constantly moves sodium out of cells and potassium into cells against the concentration gradients. During hypoxia the metabolic energy required by this pump which normally extrudes sodium from cells is lacking. As a result, sodium accumulates intracellularly. To maintain osmotic equilibrium, water shifts in, swelling the endothelial cell, impairing the flow of blood. Following prolonged depression of blood flow, tissue death results. However, if flow can be restored in time, reversal of the ischemic injury can occur. Restoration of flow, which is dependent upon the removal of intracellular water with consequent shrinking of endothelial cells to normal size, can be readily accomplished by the intravenous administration of hypertonic mannitol solutions to increase extracellular fluid osmolality.

### **Clinical Uses**

#### *Diuretic*

Being an obligatory solute, mannitol may be used either as an osmotic diuretic per se to maintain water excretion or potentiate the action of saluretic diuretics (see below). In the clinical edemas of severe nephrotic syndrome,<sup>31</sup> cirrhosis with ascites<sup>32</sup> or congestive heart failure, conventional diuretics, for example, the loop diuretics, may not promote much diuresis partly because of greatly increased proximal tubular reabsorption of sodium and water caused by ineffectively circulating volume. Under these circumstances, mannitol, by reducing proximal tubular reabsorption of sodium and water, increases the sodium load reaching the more distal tubular segments, enhancing the effect of the saluretics. Because enhanced distal nephron sodium reabsorption may quantitatively play a greater role in sodium retention in these conditions, however, the use of mannitol alone might not be sufficient to produce the desired diuresis.

Clinically, mannitol has also been used to treat severe water overload and hyponatremia. In these situations, the hyponatremia is dilutional. The osmotic effect of mannitol in the urine, as discussed earlier, causes a diuresis with an excess of water over electrolytes being excreted. By administering 25 gram boluses of hypertonic mannitol intravenously hourly, as needed to maintain urine flow, water excretion can be notably augmented. Each 5 grams of mannitol ultimately

leads to excretion of approximately 100 ml of water in excess of electrolyte, and serum sodium will rise. Thus, massive water diuresis with rapid correction of the hyponatremia follows intravenous infusion of hypertonic mannitol. However, in patients with existing intravascular volume overload or with severe myocardial disease, such infusions may excessively expand intravascular volume producing pulmonary congestion or frank pulmonary edema.

#### *Prevention of Acute Renal Failure*

Although some have questioned whether mannitol administration may prevent the development of ischemic acute renal failure (ARF),<sup>33,34</sup> considerable experimental and clinical data support this prophylactic application.<sup>35-46</sup> The reduction in ischemic injury to the kidneys of rats and dogs by prophylactic or early mannitol infusions is well documented.<sup>35-40</sup> For example, in rats mannitol infused at the time of initiation of ischemia not only increased renal blood flow<sup>36</sup> but also significantly decreased cell swelling and the resultant rise in blood urea nitrogen and creatinine.<sup>35</sup> However, in dogs, the renal failure following hemorrhagic hypotension has been reported to be unaffected by mannitol in one study,<sup>37</sup> whereas others have shown there to be significant increases in creatinine and para-aminohippurate clearances (glomerular filtration rate and renal blood flow) when mannitol was given before hypotension was induced.<sup>38-40</sup>

The benefit of mannitol in human ischemic renal disease is suggestive, if not conclusive. In 37 patients in whom oliguria developed in the absence of intravascular volume depletion, Luke and co-workers<sup>41</sup> infused 100 ml of a 20 percent mannitol solution intravenously over 10 to 30 minutes and measured urine output. If the output reached 50 ml per hour over the next two hours, no further mannitol was given. If the output increased, but less than 50 ml per hour, the same dose was repeated. By this technique, responders and nonresponders to mannitol could be identified. The responders subsequently had oliguria for less than 50 hours and urine-to-plasma osmolality ratios greater than 1.05, and renal failure did not occur. This strongly suggested that the mannitol infusion either identified the nonresponders as patients who would progress to renal failure or was able to reverse the developing renal failure in the responders as reported by Eliahou.<sup>47</sup> Similarly, in a variety of toxin or pigment induced renal

failure in animals,<sup>42,43</sup> mannitol, if given early, has been shown to ameliorate or to prevent renal failure. Finally, Dawson<sup>45,46</sup> has reported extensive experimental and clinical evidence that the severe acute tubular necrosis (ATN) which may complicate the postoperative course of hyperbilirubinemic patients may be prevented by establishing and maintaining mannitol diuresis until serum bilirubin concentration has fallen to an acceptable level.

In summary, the evidence in animals and humans is highly suggestive that mannitol, either by decreasing glomerular capillary endothelial cell swelling and thus restoring glomerular blood flow; decreasing proximal tubular cell swelling and promoting movement of intratubular debris<sup>48</sup> that could be causing tubular obstruction; diluting urine and washing out toxin; or improving renal perfusion, can ameliorate or prevent development of acute renal failure from a variety of causes. In oliguric patients with repleted intravascular volume we recommend the following mannitol trial:

- Rapidly inject 12.5 to 25 grams of 20 percent to 25 percent solution of mannitol intravenously and measure urine volume and specific gravity hourly.
- If urine output in the next hour is less than 50 ml repeat the same dose.
- If still no response, established acute renal failure is present and no more mannitol should be given.
- If the patient responds, either incipient acute renal failure has been reversed or oliguric acute renal failure has been converted to nonoliguric, either of which has a better prognosis than oliguric acute renal failure. Underlying pathogenetic factors—for example, extracellular fluid depletion—should be corrected.
- Be aware of possible complications of mannitol therapy (see below).

#### *Dialysis*

Dialysis disequilibrium is a common complication of rapid, efficient hemodialysis. Manifestations of this syndrome include nausea, vomiting, headache, blurred vision and seizures all caused by brain swelling and increased intracranial pressure. Dialysis induced hypotension (in the absence of volume depletion) and muscle cramps are also felt by many<sup>49,50</sup> to be part of this syndrome.

Most symptoms result from brain and muscle cell swelling due to water shifts following rapid removal of solute from the intravascular space with resultant disruption of the normal intracellular-extracellular fluid osmotic equilibrium. The relation of dialysis induced hypotension to osmolality changes remains to be defined. Hagstam and co-workers<sup>49</sup> and Rodrigo and associates<sup>50</sup> showed that many features of this syndrome can be prevented by infusing hypertonic mannitol into the venous return line of the dialyzer continuously during dialysis. In functionally anephric patients, however, mannitol should be used for only the first few dialyses or intermittently when severe symptoms related to disequilibrium occur<sup>50</sup> to avoid mannitol accumulation in the extracellular fluid.

Mannitol has two other possible uses in dialysis. First, it may be added to peritoneal dialysate to increase the osmolality, thus promoting fluid removal, particularly in diabetic patients who become severely hyperglycemic after continued use of conventional hypertonic glucose peritoneal dialysate. Second, mannitol infusions might be given to maintain osmotic load and urine output after dialysis preventing virtual anuria in patients undergoing hemodialysis or peritoneal dialysis therapy. Maintaining some urine flow would aid in fluid management and promote some potassium excretion in these patients. The later use for hypertonic mannitol has not been adequately tested, however.

#### *Renal Transplantation*

Mannitol may be administered safely to patients with renal transplants to distinguish post-transplant acute tubular necrosis and acute rejection. Anderson and associates<sup>51</sup> carried out studies in five renal transplant patients with previous bilateral nephrectomies, who were oliguric following transplantation. Acute rejection eventually was found in four of the five in whom diuresis occurred after injections of mannitol and ethacrynic acid.

#### *Diagnostic Tool*

Mannitol has been used to diagnose renal disease unrelated to acute renal failure. Mannitol pyelography may be used to visualize renal collecting structures not seen on routine intravenous pyelogram because of high blood urea nitrogen,<sup>52</sup> as well as to magnify the intravenous pyelogram

abnormalities associated with unilateral renal artery stenosis.<sup>50</sup> Similarly, partial versus complete and central versus renal diabetes insipidus<sup>54</sup> may be distinguished by comparing the change in free-water clearance ( $CH_2O$ ) in response to infusion of hypertonic mannitol alone (which stimulates antidiuretic hormone release and decreases free water clearance) in contrast to infusion of mannitol plus antidiuretic hormone to see if the response to mannitol can be augmented. The details are nicely outlined by Oetliker and associates.<sup>54</sup>

#### *Toxin Excretion*

Mannitol infusion has been used extensively in prophylactic and active treatment of intoxications with endogenous and exogenous substances whose clearance is increased by mannitol.<sup>55</sup> Rapid excretion of secobarbital,<sup>55</sup> eucalyptus oil,<sup>56</sup> salicylates<sup>57</sup> and bromides<sup>58</sup> has been reported after mannitol administration. In addition, endogenous substances such as uric acid,<sup>59,60</sup> hemoglobin and myoglobin are rapidly excreted without renal damage after mannitol administration. Mannitol may also modify or prevent the nephrotoxicity of certain chemotherapeutic agents. This has been clearly shown for cis-platinumdisamminedichloride<sup>61,62</sup> and is suggestive for tetracycline<sup>63</sup> and amphotericin B.<sup>64-66</sup>

#### *Cerebrovascular Disease*

Because cerebral edema or intracranial pressure is reduced following rapid infusion of 20 percent mannitol solution, this technique has been widely used in treating cerebrovascular diseases including postangiography brain swelling,<sup>67</sup> Reye syndrome,<sup>68</sup> hypoglycemia,<sup>69</sup> trauma<sup>70,71</sup> and the cerebral edema following neurosurgery,<sup>72,73</sup> as well as dialysis disequilibrium. A 20 percent to 50 percent reduction in intracranial pressure 30 to 45 minutes after mannitol administration has been noted. This effect may last up to several hours. A small rebound in pressure is then noted,<sup>74</sup> but this is much less than that seen after urea administration. An analogous effect of mannitol on spinal cord pressure after trauma in cats has been noted,<sup>75</sup> suggesting that mannitol may be used to decrease spinal cord swelling after trauma, surgical operation or irradiation. Whether mannitol can promote resorption of chronic subdural hematomas has been disputed<sup>76,77</sup> and at present is unsettled. Finally, there is a suggestion that

## MANNITOL

mannitol can alter the blood brain barrier and increase the entry of intravenously administered antibiotics into the central nervous system.<sup>78</sup>

### Cardiovascular Disease

Oliguria with decreased glomerular filtration rate and renal plasma flow commonly occurs following open heart surgical procedures under cardiopulmonary bypass. The mechanisms are not fully understood but may reflect a degree of acute renal failure. Mannitol given either intravenously or as a prime for the bypass pump, may decrease the magnitude and severity of these changes.<sup>79,80</sup> In patients in whom aortic operations are done with cross-clamping of the aorta below the renal arteries, analogous changes develop in renal function, which are also prevented by intravenous mannitol given just before cross-clamping.<sup>81</sup> Finally, more recently the protective effect

of mannitol on ischemic myocardium has been intensively studied. When given to animals just before experimental coronary artery occlusion, mannitol decreases cell swelling, water content and necrosis in anoxic myocardial cells.<sup>82-90</sup> It increases total and collateral myocardial blood flow, reduces electrocardiographic evidence of ischemia, decreases coronary vascular resistance and increases cardiac output. However, such administration of mannitol after coronary occlusion has not limited infarct size, and the role of mannitol therapy in human coronary artery disease remains to be elucidated.

### Miscellaneous Uses

Mannitol has been used to decrease pulmonary vascular resistance and increase alveolar ventilation in the acute respiratory distress syndrome,<sup>91</sup> to terminate pregnancies,<sup>92</sup> to stop hematuria as-

TABLE 1.—Clinical Applications of Intravenously Given Mannitol Solutions

<p>I. Diagnostic Evaluation of Acutely Oliguric Patient</p> <p>A. Test dose of 12.5 to 25 grams mannitol (20 percent to 25 percent solution) should be given in intravenous push over three to five minutes.</p> <p>B. If no response (&gt;50 ml urine/hour) repeat the same dose in one hour.</p> <p>C. If still no response, no further mannitol should be given and treatment for acute tubular necrosis instituted.</p>	<p>sodium chloride, 24 mEq sodium acetate, 1 gram calcium gluconate, 1 gram magnesium sulfate and 20 mEq potassium acetate. If alkalization of the extracellular fluid and urine is desired as in treatment of barbiturate intoxication less sodium chloride and more sodium acetate may be given depending on hourly checks of urine pH or blood gases. Serum electrolytes and osmolality should be monitored periodically.</p>
<p>II. Therapeutic Uses</p> <p>A. Prevention of acute renal failure (such as after cardiovascular surgical operation)</p> <ol style="list-style-type: none"> <li>1. Test dose as above followed by a priming dose of sufficient concentrated (20 percent) mannitol over an hour to bring the priming dose to 50 grams.</li> <li>2. Maintain urine output at 50 ml/hour with intravenous infusion of 5 percent mannitol plus 20 mEq sodium chloride/liter and 1 gram calcium gluconate/liter. Later, if needed, appropriate concentrations of potassium acetate, magnesium sulfate and more sodium chloride should be given as indicated.</li> </ol> <p>B. Treatment of Oliguria</p> <ol style="list-style-type: none"> <li>1. Test dose as above.</li> <li>2. If response is noted, look for underlying cause of oliguria other than acute tubular necrosis, for example, hypovolemia due to subtle cause of extracellular fluid contraction, such as third spacing.</li> </ol> <p>C. Adjunctive Therapy for Intoxications</p> <ol style="list-style-type: none"> <li>1. Test dose as above followed by priming dose (IIA).</li> <li>2. Maintain urine output at 150 to 500 ml/hour with a continuous intravenous infusion of 5 percent mannitol, each liter containing 45 mEq</li> </ol>	<p>D. Treatment of Elevated Intracranial or Intraocular Pressure</p> <ol style="list-style-type: none"> <li>1. 100 to 150 grams (15 percent or 20 percent solution) given over 30 to 60 minutes. Repeat as necessary to control intracranial or intraocular pressure.</li> <li>2. Use 60 to 90 minutes before craniotomy or ocular surgery for maximal effect.</li> </ol> <p>E. Treatment of Water Intoxication With Symptomatic Dilutional Hyponatremia</p> <ol style="list-style-type: none"> <li>1. Test dose followed by a priming dose as above (IIA).</li> <li>2. Give 20 percent mannitol as 25 gram boluses hourly or by continuous infusion at a rate of 100 to 125 ml per hour.</li> </ol> <p>F. Potentiation of Diuresis in Patients With Intractable Cardiac, Cirrhotic and Nephrotic Edema</p> <ol style="list-style-type: none"> <li>1. Give intravenous injection of near maximal dose of ethacrynic acid or furosemide.</li> <li>2. Follow immediately with test dose of mannitol as described above followed by slow intravenous infusion of 10 percent to 20 percent mannitol at a rate of 50 to 75 ml/hour.</li> <li>3. Patient's cardiovascular status including central venous pressure, urine output and osmolality and serum electrolytes should be carefully monitored during the infusion.</li> </ol>

## MANNITOL

sociated with sickle cell trait<sup>93</sup> and to augment iodine 131 uptake in the treatment of thyroid cancer.<sup>94</sup>

### Administration

Specific recommendations for intravenous administration of mannitol are presented in Table 1 for each of several clinical applications.

### Precautions

One case report of an allergic reaction associated with mannitol administration has been published.<sup>95</sup> However, the relationship is unclear since mannitol is inert and no other cases have been reported. On the other hand, in a patient with poor renal function and incipient heart failure, excessive volume expansion following mannitol infusion can be a serious side effect and mannitol should be administered very cautiously in such a situation. After hypertonic mannitol administration, particularly of bolus doses,<sup>96</sup> dilutional hyponatremia and cellular dehydration may result from the movement of water out of cells in response to increased extracellular fluid tonicity. In a patient with good renal function this abnormality is of little consequence and is rapidly corrected as mannitol is excreted. In a patient with poor kidney function, however, sustained hyperosmolality, hyponatremia and cellular dehydration may be serious complications.<sup>97</sup> Rarely, hypernatremia and dehydration from excessive water loss in response to larger rapid infusions of hypertonic mannitol in patients with functioning kidneys, and occasionally mild hyperkalemia of unclear origin,<sup>98</sup> may occur. Finally, in uremic dogs given mannitol pronounced decreases in cerebrospinal fluid pH resulting in death have been noted.<sup>99</sup> Similar changes in cerebrospinal fluid pH in human beings with uremia after mannitol administration have not been studied.

### REFERENCES

- Smith WW, Finkelstein N, Smith HW: Renal excretion of hexitols and their derivatives and of endogenous creatinine-like chromogen in dog and man. *J Biol Chem* 135:231-250, 1940
- Oldendorf WH, Kitano M: The early disappearance of extracellular tracers from plasma after I.V. injection. *Soc Exp Biol Med* 141:940-943, 1972
- Dominguez R, Corcoran AC, Page IH: Mannitol—Kinetics of distribution, excretion and utilization in human beings. *J Lab Clin Med* 32:1192-1202, 1947
- Wesson LG, Anslow P: Excretion of sodium and water during osmotic diuresis in dogs. *Am J Physiol* 153:465-474, 1948
- Seely JF, Dirks JH: Micropuncture study of hypertonic mannitol diuresis in the proximal and distal tubule of the dog kidney. *J Clin Invest* 48:2330-2340, 1969
- Gennari FJ, Kassirer JP: Osmotic diuresis. *N Engl J Med* 291:714-720, 1974
- Lilien OM, Jones SG, Mueller CB: The mechanism of mannitol diuresis. *Surg Gynecol Obstet* 117:221-228, 1963
- Ternes SP, Lilien OM, Chamberlain W: A direct vasoconstrictor effect of mannitol on the renal artery. *Surg Gynecol Obstet* 141:223-226, 1975
- B'Alint P, Szocs E, Tari'an E, et al: Autoregulation of renal circulation in mannitol-loaded dogs. *Int Urol Nephrol* 7:65-78, 1975
- Velasquez MT, Notargiacomo AV, Cohn JN: Comparative effects of saline and mannitol on renal cortical blood flow and volume in the dog. *Am J Physiol* 224:322-327, 1973
- Detmer DE, Zimmerman JM, King TC: Mannitol diuresis: The relationship of plasma volume to renal blood flow. *J Surg Res* 5:552-555, 1965
- Lilien CM: The paradoxical reaction of renal vasculature to mannitol. *Invest Urol* 10:346-353, 1973
- Parfitt AM: The acute effects of mersalyl, chlorothiazide and mannitol on the renal excretion of calcium and other ions in man. *Clin Sci* 36:267-282, 1969
- Sotorn'ik I, Schuck O, Stribrna J: Influence of diuretics on renal calcium excretion. *Experientia* 25:591-592, 1969
- Maesaka JK, Berger ML, Bornia ME, et al: Effect of mannitol on phosphate transport in intact and acutely thyroparathyroidectomized rats. *J Lab Clin Med* 87:680-691, 1976
- DiScala VA, Mautner W, Cohen JA, et al: Tubular alterations produced by osmotic diuresis with mannitol. *Ann Intern Med* 63:767-775, 1965
- Janssen CW Jr: Osmotic nephrosis—A clinical and experimental investigation. *Acta Chir Scand* 134:481-487, 1968
- Barry KG, Berman AR: Mannitol infusion—III. The acute effect of the intravenous infusion of mannitol on blood and plasma volumes. *N Engl J Med* 264:1085-1088, 1961
- Hoff JE, Deavers S, Huggins RA: Effect of hypertonic glucose and mannitol on plasma volume. *Proc Soc Exp Biol Med* 122:630-634, 1966
- Willerson JT, Curry GC, Atkins JM, et al: Influence of hypertonic mannitol on ventricular performance and coronary blood flow in patients. *Circulation* 51:1095-1100, 1975
- Atkins JM, Wildenthal K, Horwitz LD: Cardiovascular responses to hyperosmotic mannitol in anesthetized and conscious dogs. *Am J Physiol* 225:132-137, 1973
- Raizner AE, Costin JC, Croke RP, et al: Reflex, systemic and local hemodynamic alterations with experimental hyperosmolality. *Am J Physiol* 224:1327-1333, 1973
- Krishnamurthy VS, Adams HR, Smitherman TC, et al: Influence of mannitol on contractile responses of isolated perfused arteries. *Am J Physiol* 232:H59-H66, 1977
- Goluboff B, Shenkin HA, Haft H: The effects of mannitol and urea on cerebral hemodynamics and cerebrospinal fluid pressure. *Neurology* 14:891-898, 1964
- Cottrell JE, Robustelli A, Post K, et al: Furosemide and mannitol induced changes in intracranial pressure and serum osmolality and electrolytes. *Anesthesiology* 47:28-30, 1977
- Adams RE, Kirschner RJ, Leopold IH: Ocular hypotensive effect of intravenous mannitol. *Arch Ophthalmol* 69:55-58, 1963
- Zayed A, Fouad AR: The role of mannitol osmotherapy in retinal detachment surgery. *Bull Ophthalmol Soc Egypt* 64:483-490, 1971
- Lazar M, Nowakowski J, Furman M, et al: Ocular penetration of topically applied mannitol. *Am J Ophthalmol* 70:849-851, 1970
- Ames A, Wright RL, Kowada M, et al: Cerebral ischemia—II. The no-reflow phenomenon. *Am J Pathol* 52:437-453, 1968
- Flores JE, Frega NS, DiBona DR, et al: The role of cell swelling in ischemic renal injury. In Friedman EA, Eliahou HE (Eds): *Proc Conf Renal Failure*. DHEW Publication Number (NIH) 74-608. New York, 1973, pp 19-29
- Bernstein LM, Grossman A: Diuretic effect of mannitol in nephrotic edema. *J Lab Clin Med* 59:309-319, 1962
- Gagnon O, Gertman PM, Iber FL: Comparison of oral isosorbide and intravenous mannitol as adjuncts to the diuresis of patients with cirrhosis of the liver. *Am J Med Sci* 254:284-295, 1967
- Oken DE: Mannitol and the prevention of vasomotor nephropathy. In Giovannetti S, et al (Eds): *Sixth International Congress of Nephrology*. Basel, Karger, 1976, pp 578-583
- Schroder K, Gassler U: The influence of mannitol, ethacrynic acid and furosemide on glomerular filtration in experimental acute renal failure. *Postgrad Med J* 47(Suppl):11-12, 1971
- Flores J, DiBona DR, Beck CH, et al: The role of cell swelling in ischemic renal damage and the protective effect of hypertonic solute. *J Clin Invest* 51:118-126, 1972
- Franklin WA, Ganote CE, Jennings RB: Blood reflow after renal ischemia—Effects of hypertonic mannitol on reflow and tubular necrosis after transient ischemia in the rat. *Arch Pathol* 98:106-111, 1974
- Danielson BG, Grangsjö G, Karkmark B, et al: Kidney function and intrarenal blood flow distribution after bleeding and infusions of mannitol and dextran. *Acta Aesthesiol Scand* 17:8-21, 1973
- Conte NF, Gagnon JA, Barry KG: Intrinsic renal effect of hypertonic mannitol infusion (HMI) on renal clearances of PAH ( $C_{PAH}$ ), exogenous creatinine ( $C_{Cr}$ ), and urine flow (UF) during hemorrhagic shock in dogs. *Clin Res* 10:247, 1962

# MANNITOL

39. Szabo G, Magyar S: Effect of hypertonic mannitol solution on circulation and renal function in acute blood loss. *Acta Med Acad Sci Hung* 23:325-335, 1967
40. Doberneck RC, Schwartz FD, Barry KG: A comparison of the prophylactic value of 20 percent mannitol, 4 percent urea, and 5 percent dextrose on the effects of renal ischemia. *J Urol* 89:300-302, 1963
41. Luke RG, Briggs JD, Allison ME, et al: Factors determining response to mannitol in acute renal failure. *Am J Med Sci* 259:168-174, 1970
42. Sherwood T, Lavender JP, Russell SB: Mercury-induced renal vascular shut-down—Observations in experimental acute renal failure. *Eur J Clin Invest* 4:1-8, 1974
43. Wilson DR, Thiel G, Arce ML, et al: Glycerol induced hemoglobinuric acute renal failure in the rat—3. Micropuncture study of the effects of mannitol and isotonic saline on individual nephron function. *Nephron* 4:337-355, 1967
44. Powers SR, Boba A, Hostnik W, et al: Prevention of post-operative acute renal failure with mannitol in 100 cases. *Surgery* 55:15-23, 1964
45. Dawson J: Jaundice and anoxic renal damage—Protective effect of mannitol. *Br Med J* 1:810-811, 1964
46. Dawson J: Post-operative renal function in obstructive jaundice: Effect of a mannitol diuresis. *Br Med J* 1:82-86, 1965
47. Eliahou H: Mannitol therapy in oliguria of acute onset. *Br Med J* 1:807-809, 1964
48. Frega NS, DiBona DR, Leaf A: Ischemic renal injury. *Kidney Int* 10:517-525, 1976
49. Hagstam KE, Lindergard B, Tibbling G: Mannitol infusion in regular haemodialysis treatment for chronic renal insufficiency. *Scand J Urol Nephrol* 3:257-263, 1969
50. Rodrigo F, Shideman J, McHugh R, et al: Osmolality changes during hemodialysis. *Ann Intern Med* 86:554-561, 1977
51. Anderson CF, O'Kane HO, Shorter RG, et al: Use of diuretic agents during oliguria after renal transplantation. *Surgery* 67:249-254, 1970
52. Taylor DA, Macken KL, Fiore AS: Mannitol pyelography: A simplification of the drip infusion technic. *Radiology* 88:1117-1120, 1967
53. Taylor DA, Chovnick SD, Phillips SH: The use of rapid sequence mannitol pyelography in the diagnosis of renovascular hypertension. *Radiology* 89:1095-1099, 1967
54. Oetliker OH, Simon J, Tietze HU: Diagnostic value of mannitol-induced diuresis in children. *Acta Paediatr Scand* 63:113-121, 1974
55. Cirksema WJ, Bastain RC, Malloy JP, et al: Use of mannitol in exogenous and endogenous intoxications. *N Engl J Med* 270:161-166, 1964
56. Gurr FW, Scroggie JE: Eucalyptus oil poisoning treated by dialysis and mannitol infusion, with an appendix on the analysis of the biological fluids for alcohol and eucalyptol. *Australia Ann Med* 14:238-249, 1965
57. Prowse K, Pain M, Marston AD, et al: The treatment of salicylate poisoning using mannitol and forced alkaline diuresis. *Clin Sci* 38:327-337, 1970
58. Adamson JS Jr, Flanagan WJ, Ackerman GL: Treatment of bromide intoxication with ethacrynic acid and mannitol diuresis. *Ann Intern Med* 65:749-752, 1966
59. Barry KG, Hunter RH, David TE, et al: Acute uric acid nephropathy: Treatment with mannitol diuresis and peritoneal dialysis. *Arch Intern Med* 111:452-459, 1963
60. Skeith MD, Healey LA, Cutler RE: Urate excretion during mannitol and glucose diuresis. *J Lab Clin Med* 70:213-220, 1967
61. Cvitkovic E, Spaulding J, Bethune V, et al: Improvement of cis-dichlorodiammine-platinum (NSC 119875): Therapeutic index in an animal model. *Cancer* 39:1357-1361, 1977
62. Hayes DM, Cvitkovic E, Golbey RB, et al: High dose cis-platinum diammine dichloride: Amelioration of renal toxicity by mannitol diuresis. *Cancer* 39:1372-1381, 1977
63. Polec RB, Yeh SD, Shils ME: Protective effect of ascorbic acid, isoascorbic acid and mannitol against tetracycline-induced nephrotoxicity. *J Pharmacol Exp Ther* 178:152-158, 1971
64. Olivero JJ, Lozano-Mendez J, Ghafary EM, et al: Mitigation of amphotericin B nephrotoxicity by mannitol. *Br Med J* 1:550-551, 1975
65. Bullock WE, Luke RG, Nuttall CE, et al: Can mannitol reduce amphotericin B nephrotoxicity?—Double-blind study and description of a new vascular lesion in kidneys. *Antimicrob Agents Chemother* 10:555-563, 1976
66. Rosch JM, Pazin GJ, Fireman P: Reduction of amphotericin B nephrotoxicity with mannitol. *JAMA* 235:1995-1996, 1976
67. Bouzarth WF, Goldfeder P, Shenkin HA: Hypertonic mannitol as a treatment for complications of cerebral arteriography. *Am J Roentgen* 104:119-122, 1968
68. Kindt GW, Waldman J, Kohl S, et al: Intracranial pressure in Reye syndrome—Monitoring and control. *JAMA* 231:822-825, 1975
69. MacCulsh AC, Munro JF, Duncan LJ: Treatment of hypoglycaemic coma with glucagon intravenous dextrose, and mannitol infusion in a hundred diabetics. *Lancet* 2:946-949, 1970
70. Parker AJ, Park RD, Stowater JL: Reduction of trauma-induced edema of spinal cord in dogs given mannitol. *Am J Vet Res* 34:1355-1357, 1973
71. Shalit MN: Effect of intracarotid artery administration of mannitol on cerebral blood flow and intracranial pressure in experimental brain edema. *Isr J Med Sci* 10:577-580, 1974
72. Wise BL, Chater N: The value of hypertonic mannitol solution in decreasing brain mass and lowering cerebral spinal fluid pressure. *J Neurosurg* 19:1038-1043, 1962
73. Shenkin HA, Goluboff B, Haft H: The use of mannitol for the reduction of intracranial pressure in intracranial surgery. *J Neurosurg* 19:897-901, 1962
74. Shenkin HA, Goluboff B, Haft H: Further observations on effects of abnormally increased osmotic pressure of plasma on cerebrospinal-fluid pressure in man. *J Neurosurg* 22:563-568, 1965
75. Richardson HD, Nakamura S: An electron microscopic study of spinal cord edema and the effect of treatment with steroids, mannitol, and hypothermia. *Proc Vet Adm Spinal Cord Inj Conf* 18:10-16, 1971
76. Gjerris F, Schmidt K: Chronic subdural hematoma—Surgery or mannitol treatment. *J Neurosurg* 40:639-642, 1974
77. Suzuki J, Takaku A: Nonsurgical treatment of chronic subdural hematoma. *J Neurosurg* 33:548-553, 1970
78. Takimoto N, Watanabe M, Kinjo T, et al: Clinical study on penetration of antibiotics into cerebrospinal fluid (second report)—Study on intravenous administration of sodium cephalothin with mannitol solution. *Neuro Surg (Tokyo)* 5:137-143, 1977
79. Kahn DR, Cerny JC, Lee RWS, et al: The effect of dextran and mannitol on renal function during open-heart surgery. *Surgery* 57:676-679, 1965
80. Etherige EE, Levitin H, Nakamura K, et al: Effect of mannitol on renal function during open-heart surgery. *Ann Surg* 161:53-62, 1965
81. Barry KG, Cohen A, LeBlanc PC Jr: Mannitolization—I. The prevention and therapy of oliguria associated with cross-clamping of the abdominal aorta. *Surgery* 50:335-340, 1961
82. Powell WJ Jr, DiBona DR, Flores J, et al: Effects of hyperosmotic mannitol in reducing ischemic cell swelling and minimizing myocardial necrosis. *Circulation* 53:45-49, 1976
83. Powell WJ Jr, DiBona DR, Flores J, et al: The protective effect of hyperosmotic mannitol in myocardial ischemia and necrosis. *Circulation* 54:603-615, 1976
84. Kloner RA, Reimer KA, Willerson T, et al: Reduction of experimental myocardial infarct size with hyperosmolar mannitol. *Proc Soc Exp Biol Med* 151:677-683, 1976
85. Hutton I, Curry GC, Templeton GH, et al: Influence of hypertonic mannitol on regional myocardial blood flow and ventricular performance in awake, intact dogs with prolonged coronary artery occlusion. *Cardiovascular Res* 9:409-419, 1975
86. Willerson JT, Weisfeldt ML, Sanders CA, et al: Influence of hyperosmolar agents on hypoxic cat papillary muscle function. *Cardiovascular Res* 8:8-17, 1974
87. Christodoulou J, Erlandson R, Smithen C, et al: Effects of mannitol on cardiac ultrastructure and microcirculation following anoxia. *Am J Physiol* 229:853-860, 1975
88. Smithen C, Christodoulou J, Killip T, et al: Metabolic and hemodynamic consequences of mannitol following myocardial anoxia. *Am J Physiol* 229:847-852, 1975
89. Brachfeld N: Metabolic evaluation of agents designed to protect the ischemic myocardium and to reduce infarct size. *Am J Cardiol* 37:528-532, 1976
90. Brachfeld N, Christodoulou J, Keller N, et al: Hemodynamic, metabolic, and ultrastructural consequences of hyperosmolar mannitol after myocardial anoxia. *Recent Adv Studies Cardiac Struct Metab* 7:459-466, 1975
91. Powers SR Jr, Dhiraj S, Ryon D, et al: Hypertonic mannitol in the therapy of the acute respiratory distress syndrome. *Ann Surg* 185:619-625, 1977
92. Deshmukh MA, Shah S, Shah S, et al: Extra-amniotic mannitol for pregnancy termination (preliminary report). *J Postgrad Med* 22:23-25, 1976
93. Knochel JP: Hematuria in sickle cell trait: The effect of intravenous administration of distilled water, urinary alkalization, and diuresis. *Arch Intern Med* 123:160-165, 1969
94. Hamburger JJ, Desai P: Mannitol augmentation of I-131 uptake in the treatment of thyroid carcinoma. *Metabolism* 15:1055-1058, 1966
95. Spaeth GL, Spaeth EB, Spaeth PG, et al: Anaphylactic reaction to mannitol. *Arch Ophthalmol* 78:583-584, 1967
96. Tunner WS, Kiser WS, Mann P: Studies on the significance of decreased serum sodium levels following the clinical use of mannitol. *J Urol* 94:470-474, 1965
97. Aviram A, Prau A, Czaczkes JW, et al: Hyperosmolality with hyponatremia, caused by inappropriate administration of mannitol. *Am J Med* 42:648-650, 1967
98. Moreno M, Murphy C, Goldsmith C: Increase in serum potassium resulting from the administration of hypertonic mannitol and other solutions. *J Lab Clin Med* 73:291-298, 1969
99. Silber SJ, Thompson N: Mannitol induced central nervous system toxicity in renal failure. *Invest Urol* 9:310-312, 1972